AUTHOR INDEX

Alton, G. 339 Axelos, M.A.V. 271

Baer, H.H. 215
Basora, N. 1
Beaupere, D. 327
Beri, R.G. 11
Brade, H. 93
Brade, L. 93
Brazier, M. 327
Brimacombe, J.S. 313, 317
Brisson, J.-R. 49
Buchholz, K. 307

Čadová, E. 321 Čapek, K. 321 Cerezo, A.S. 241 Chittenden, G.J.F. 185 Chornet, E. 1 Ciancia, M. 241 Crépeau, M.-J. 271

Demailly, G. 327

Eyrisch, O. 287

Fessner, W.-D. 287 Finch, P. 241 Fujita, K. 75 Fukase, K. 93

Gonsho, A. 163

Hara, K. 75 Hindsgaul, O. 339

Irie, K. 163 Ito, Y. 163 Iwasawa, H. 163

Jann, B. 261, 335 Jann, K. 261, 335 Kamerling, J.P. 135 Kanie, Y. 339 Kerékgyártó, J. 135 Kitahata, S. 75 Kogan, G. 261, 335 Koizumi, K. 75 Köll, P. 313, 317 Kondo, T. 231 Kopf, J. 313, 317 Kosma, P. 93 Kotani, K. 109, 193 Koyama, Y. 147 Kusumoto, S. 93 Kuwahara, N. 75

Lieker, H.-P. 307 Lipták, A. 135

März, L. 93 Matulewicz, M.C. 241 Meyer, B. 49 Morf, M. 313, 317

Niola, F. 1 Nogusa, H. 163

Oguma, T. 193

Peters, T. 49

Raaijmakers, H.W.C. 185 Reese, E.T. 11 Reilly, P.J. 307 Renard, C.M.G.C. 271 Rochas, C. 27 Roger, P. 271 Rollings, J.E. 11 Roos, Y. 39

Saito, K. 109, 193 Sedmera, P. 321 Shashkov, A.S. 261 Shibukawa, M. 163 Sinerius, G. 287 Somorjai, R. 49 Stephen, A.M. 249 Strobl, M. 93 Stuike-Prill, R. 49 Sugawara, T. 163 Susaki, H. 163

Tanimoto, T. 75 Taravel, F.-R. 27 Thibault, J.-F. 271 Thielecke, K. 307 Tobe, K. 193 Tokutake, S. 109, 193 Tsuchiya, T. 147 Turquois, T. 27

Uchida, R. 109 Umemura, E. 147 Umezawa, S. 147 Uzan, R. 327

Van der Ven, J.G.M. 135 Vidal, P.F. 1 Vliegenthart, J.F.G. 135 Vogt, D.C. 249

Wadouachi, A. 327 Walker, J. 11 Wu, X. 215

Yamaji, N. 109, 193

Zimmer, B. 313, 317 Zwanenburg, B. 185

SUBJECT INDEX

- N-Acetylation, determination of chitin-chitosan samples of the degree of, 1
- α-N-Acetylneuraminic acid, structure of the polysaccharide component of the O56 antigen of Escherichia coli which contains 7-substituted, 261
- Acceptor for N-acetylglucosaminyltransferase-III, use of a synthetic dideoxygenated pentasaccharide as a specific, 339
- 2,3-Anhydro-4,6-O-benzylidene-α-p-allopyranoside, an alternative synthesis of methyl, using carbonate esters, 185
- 1,3-Anhydro-β-D-xylo-hexulofuranoside, α-D-glucopyranosyl, an intermediate in the alkaline hydrolysis of α-D-glucopyranosyl 3,4-anhydro-β-D-lyxo-hexulofuranoside, 321
- 6-O-Alkylcelluloses, preparation, 231
- Alkaline hydrolysis of α-D-glucopyranosyl 3,4anhydro-β-D-dyxo-hexulofuranoside, formation of α-D-glucopyranosyl 1,3-anhydro-β-Dxylo-hexulofuranoside as an intermediate in the, 321
- Amylase, synthesis of modified 2-chloro-4nitrophenyl β-maltopentaosides as useful substrates for assay of human alpha, 109
- Amylases, syntheses of subtractively modified 2-chloro-4-nitrophenyl β-maltopentaosides, and their application to the differential assay of human, alpha-, 193
- Antigen of Escherichia coli, structure of the neuraminic acid-containing O24 polysaccharide, 335
- Antigen of Escherichia coli, structure of the O56 polysaccharide that contains 7-substituted α-N-acetylneuraminic acid, 261
- Building block that is part of xylose-containing carbohydrate chains from N-glycoproteins, synthesis of a selectively protected trisaccharide, 135
- Carbonate esters, an alternative synthesis of methyl 2,3-anhydro-4,6-O-benzylidene-α-Dallopyranoside using, 185

- Carrageenans from Gigartina skottsbergii, determination of the structures of cystocarpic, by methylation analysis and NMR spectroscopy, 241
- Celluloses, 6-O-alkyl (ethyl, methyl), preparation, 231
- Chitin, determination of the degree of N-acetylation of, 1
- Chitosan, determination of the degree of Nacetylation of, 1
- Chitosan via coupled size-exclusion chromatography and multiple-angle laser light-scattering technique, characterization of, 16
- Conformational analysis of saccharides using a Monte Carlo method, 49
- Copolymers of poly(acrylamide) and modified Kdo residues, synthesis and immunoreactivity of, 93
- Crystal and molecular structure of deca-Oacetyl-L-galacto-L-gulo-decitol, 313
- Crystal and molecular structure of octa-Oacetyl-L-threo-D-galacto-octitol, 317
- Cyclomalto-oligosaccharides having β-D-galactosyl residue(s) on the side chain, preparation, isolation, and characterization of novel heterogeneous branched, 75
- Decitol, crystal and molecular structure of deca-O-acetyl-L-galacto-L-gulo-, 313
- Dideoxygenated pentasaccharide as a specific acceptor for N-acetylglucosaminyltransferase-III, use of a synthetic, 339
- Encephalartos longifolius, further hydrolytic studies of the gum exudate of, 249
- Enzymic de novo synthesis of D-tagatose 1,6-bisphosphate, 287
- Fluoro- and 5-deoxy-5,5-difluoro-netilmicin, synthesis of 5- deoxy-5-, 147
- Galacturonic regions in pectins, studies by acid hydrolysis of the length of homo-, 271

3

α-D-Glucopyranosyl 1,3-anhydro-β-D-xylo-hexulofuranoside, an intermediate in the alkaline hydrolysis of α-D-glucopyranosyl 3,4anhydro-β-D-bxo-hexulofuranoside, 321

D-Glucose derivatives, esterification of, by oxolinic acid, 327

Glycopyranosides and of their conjugates with poly(L-glutamic acid) having a 2-(4-hydroxyphenyl)ethylamino group for radiolabeling, synthesis of 8-aminoctyl, 163

Gum exudate of Encephalartos longifolius, further hydrolytic studies, 249

High-performance anion-exchange chromatography of saturated and unsaturated oligogalacturonic acids, 307

Homogalacturonic regions in pectins, studies of the length by acid hydrolysis, 271

Hydrolytic studies of the gum exudate of Encephalartos longifolius, further, 249

Kdo residues, modified, synthesis and immunoreactivity of poly(acrylamide) copolymers containing, 93

Light-scattering technique, characterization of chitosan via coupled size-exclusion chromatography and multiple-angle laser, 16

Low molecular weight carbohydrates, melting and glass transitions of, 39

β-Maltopentaosides as useful substrates for assay of human alpha amylase, synthesis of modified 2-chloro-4-nitrophenyl, 109

β-Maltopentaosides, syntheses of subtractively modified 2-chloro-4-nitrophenyl, and their application to the differential assay of human alpha-amylases, 193

Melting and glass transitions of low molecular weight carbohydrates, 39

Methyl 2,3-anhydro-4,6-O-benzylidene-α-D-allopyranoside, an alternative synthesis of methyl, using carbonate esters, 185

Methylation analysis and NMR spectroscopy, determination of the structures of cystocarpic carrageenans from Gigartina skottsbergii by, 241

Molecular and crystal structure of deca-Oacetyl-L-galacto-L-galo-decitol, 313

Molecular and crystal structure of octa-Oacetyl-1-threo-D-galacto-octitol, 317

Monte Carlo method for conformational analysis of saccharides, 49

Netilmicin, synthesis of 5-deoxy-5-fluoro- and 5-deoxy-5,5- difluoro-, 147 Neuraminic acid-containing polysaccharide, structure of the O24 antigen of Escherichia coli, 335

NMR and rheological studies of the agarosecarob galactomannan blend, 27

NMR spectroscopy and methylation analysis, determination of the structures of cystocarpic carrageenans from Gigartina skottsbergii by, 241

Octitol, crystal and molecular structure of octa-O-acetyl-L-threo-D-galacto-, 317

Oligogalacturonic acids, saturated and unsaturated, high-performance anion-exchange chromatography of, 307

Oxolinic acid, esterification of p-glucose derivatives by, 327

Palmitic acid, 2,3- and 2,3'-diesters with α , α trehalose, synthesis, 215

Pectins, studies of the length of homogalacturonic regions by acid hydrolysis, 271

Poly(acrylamide) copolymers containing modified Kdo residues, 93

Poly(L-glutamic acid) having a 2-(4-hydroxyphenyl)ethylamino group for radiolabeling, synthesis of 8-aminoctyl glycopyranosides and of their conjugates with, 163

Polysaccharide containing neuraminic acid, structure of the O24 antigen of Escherichia

Polysaccharide containing 7-substituted α-Nacetylneuraminic acid, a component of the O56 antigen of Escherichia coli, 261

Size-exclusion chromatography and multiple-angle laser light-scattering technique, characterization of chitosan via coupled, 16

Stearic acid, 2,3- and 2,3'-diesters with α,α trehalose, synthesis, 215

Subtractively modified 2-chloro-4-nitrophenyl βmaltopentaosides and their application to the differential assay of human alphaamylases, syntheses of, 193

Synergy of the agarose carob galactomannan blend, 27

Synthesis of 8-aminoctyl glycopyranosides and of their conjugates with poly(L-glutamic acid) having a 2-(4-hydroxyphenyl)ethylamino group for radiolabeling, 163

Synthesis of 5-deoxy-5-fluoro- and 5-deoxy-5,5difluoro-netilmicin, 147

Synthesis of modified 2-chloro-4-nitrophenyl βmaltopentaosides as useful substrates for assay of human alpha amylase, 109

- D-Tagatose 1,6-bisphosphate, facile enzymic de novo synthesis and NMR spectroscopic characterization, 287
- Transitions of low molecular weight carbohydrates, melting and glass, 39
- α,α-Trehalose 2,3- and 2,3'-diesters with palmitic and stearic acid: potential immunoreactants for the serodiagnosis of tuberculosis, synthesis of, 215
- Trisaccharides building block that is part of
- xylose-containing carbohydrate chains from N-glycoproteins, synthesis of a selectively protected, 135
- Tuberculosis, potential immunoreactants for the serodiagnosis of, 215
- Xylose-containing carbohydrate chains from Nglycoproteins, synthesis of a selectively protected trisaccharide building block that is part of, 135